

## Body Mass Index (BMI) and Anthropometric Measurement of the Developing Foetus (Volume 3, Issue 1, July- September 2019 )

Author: Prof Niranjan Bhattacharya

Anthropometry is a technique implemented for the scientific study of the measurements of the human body. Over the years anthropometric measurements have become an important tool for clinicians and scientists in health assessment of the foetal development and growth<sup>1</sup>. One of the main parameters used in anthropometry to study the impact of the foetal growth through different phases of pregnancy include that of the parental and the foetal body mass index (BMI) which is believed to affect the baby's growth, development, and birth weight<sup>2</sup>.

Birth weight is also one of the important factors affecting the foetal anthropometric measurements and any abnormality is strongly associated with the morbidity and even mortality of the infant. A low birth weight of a neonate can be linked to impaired overall growth and development of the baby and overweight can be related to severe complications of the child during delivery<sup>3</sup>. Foetal and offspring overweight can be also related to obesity in children in later life<sup>3,4</sup>.

Apart from the weight of the neonate, factors such as the maternal height, weight, metabolic rate are also important anthropometric parameters related to the growth and development of the foetus<sup>5</sup>. Small and selected studies<sup>6-11</sup> have suggested that pre-pregnancy BMI (pre-BMI) and weight during the gestation period of the mother are important factors that predict the outcome of the foetal weight and its development.

Normally it is gestational weight, which is indicative of both nutritional status and tissue development whereas pre-BMI only reflects the nutritional status<sup>12</sup>. However, the relationship between the two is still not clearly established including that of the maternal BMI and other anthropometric parameters related to the foetal growth and development in pregnancy<sup>2</sup>.

An early pioneering study<sup>13</sup> involving anthropometric measures of the aborted human foetus collected ethically from all the trimesters was conducted by Mukherjee *et al.* They found the weight of the liver decreased with increase in gestational period whereas organs such as lungs increased in their size with increase in the gestational period<sup>13,14</sup>. The adrenal glands in the first trimester remain larger than the kidneys, however, after 12 weeks of gestation the kidney outweighs the adrenal gland.

The group was unable to detect any thymus tissue at 8 weeks of gestation in smaller foetuses. However, they could detect the presence of the thymus gland in larger foetuses weighing more than 5g between the first and second trimesters' period. In 28 weeks old foetuses, the thymus could be easily detected indicating that with an increase in the gestational week, the thymus becomes observable because of its increase in weight and size<sup>15</sup>. In case of sexual organs, the growth and development of the testes were not uniform. However, a trend was observed where there was a decrease in the weight of the testes with an increase in the gestational period

in the male foetuses and a similar pattern was also observed in case of the ovaries in female foetuses<sup>13</sup>.

### **Relationship between the parental BMI and the anthropometric development of the foetus :**

Most of the studies conducted are based on the relationship of birth weight as an important anthropometric parameter and its measure of the foetal growth and development<sup>16</sup>. Ay *et al*<sup>2</sup>, for the first time examined the relationship of maternal anthropometrics with the foetal growth and development at various stages of pregnancy through a large cohort study. They showed that maternal pre-BMI gestational weight and height has an influence on the foetal growth and development starting from mid-pregnancy onwards. These maternal parameters can be associated with the small and large size of the infants and are related to an increasing gestational age. The findings of the study were independent of the social factors such as the socio-economic condition and lifestyle of the mother. Studies pertaining to the timing of the anthropometric development of the foetus have been focused mainly on the birth weight and foetal growth. However, third-trimester studies till date remains inconclusive<sup>17-21</sup>.

Factors such as diabetes or insulin resistance have a strong effect on the high maternal pre-BMI and weight gain and can lead to increased foetal glucose and an increased birth weight along with a risk of cardiometabolic disorder<sup>21</sup>. Maternal nutritional status is also an important criterion that can be strongly associated with the pre-BMI and the outcome of the fetal weight<sup>22</sup>. However, further studies are important to prove this exclusively including the maternal anthropometric mechanisms which affect the foetal growth<sup>2</sup>.

Another important study<sup>23</sup> was conducted in Pune to find out the relationship of BMI and height on 557 pregnancies with foetal age between 17 and 29 weeks of gestation and observed through ultrasound method. The group reported that parental height was positively associated with an increase in the foetal head circumference and femur length. In case of higher BMI rates in mothers, ultrasound images showed that the foetus in utero had a smaller head circumference at 17 weeks and it increased during the time of birth. In mothers having lower BMI rates, head to foetal body ratio was observed to be large at 17 weeks. The placental volume was also depended on the maternal BMI and the paternal height<sup>23</sup>.

As mentioned before, pre-BMI is an important parameter and marker for nutrition, energy and tissue development<sup>23</sup>. In the above study, further, the group of clinicians found a positive correlation between the paternal height and placental volume at around 17 weeks of foetal development although the direct role of maternal BMI on the 17 weeks' foetus was found to be negative. Better maternal nutrition is thought to provide the mother with a higher BMI which helps in the placental development in early pregnancy resulting in a greater foetal development in the final gestational phases<sup>24</sup>. Similarly like Goldenberg *et al*<sup>25</sup>, the study further found a positive relationship with head circumference growth in the 29 weeks to birth interval.

A study by Zoya *et al*<sup>26</sup>, showed a weak association of maternal BMI with respect to the sex of the first-trimester foetus. They showed maternal BMI was related to the female foetus rather than the male foetus and may be sex-dependent in nature. Other factors that can influence the impact of maternal BMI on birth weight of the foetus include maternal age, ethnicity, gestational diabetes mellitus and insulin resistance, education and environmental factors, hypertension including genetics<sup>27,28</sup>.

Until recently the relationship of BMI and its significant influence on the developing foetus was unknown. Studies<sup>2,29-31</sup> and observations have come to acknowledge the fact that maternal weight, BMI including that of the foetus plays an important role in determining the foetal growth and development.

Recently the significance of paternal BMI and dietary behaviour in animal studies have also shed light into the fact that paternal obesity including BMI can affect the offspring in a gender dependant manner<sup>32</sup>. Chen *et al*<sup>32</sup>, in one such pioneer study observed that paternal BMI during the time of conception can be sex dependent and can influence the male but not the female foetus through mechanisms hitherto unknown. However, the above cannot be exclusively confirmed as of yet due to very few studies relating to paternal BMI and its relationship with the foetal development<sup>33</sup>.

#### **Importance of BMI in foetal growth:**

BMI is one of the important parameters used along with other measures such as waist circumference, waist to hip and height ratio, subcapsular thickness as a part of the classification of overweight, obesity although it fails to account for the overall fat distribution. It is also an important predictor to assess the healthy outcome of a baby. BMI can be directly correlated to neonate obesity, preterm birth complications, shoulder dystocia and other complications<sup>34</sup>. Nonetheless BMI still is an important factor that influences the growth and development of the foetus on a more general basis<sup>35</sup>.

In third world countries where maternal nutrition has a profound effect on the foetal growth and development during pregnancy, the anthropometric parameter such as intra-uterine growth (IUG) chart alone is not enough to assess the level of foetal malnutrition. BMI which is based on weight to length ratio is further an effective and sensitive method to assess the level of malnutrition<sup>36</sup>. As pregnancy is an important period for the foetal growth and development, any metabolic changes like maternal weight gain due to diabetes, dyslipidaemia or weight loss due to chronic infections and diseases can affect the foetal health and development. Foetal macrosomia, limited or stunted growth of the foetus, pre-term delivery of the offspring are often the result of maternal pre-BMI dysregulation<sup>37</sup>. Hence, maternal pre-BMI can be an effective tool in poor countries to anticipate and predict neonatal health complications apart from nutritional care<sup>37</sup>.

Glucose being one of the major energy substrates, has shown to cross the placenta *via* facilitated diffusion mechanism but no evidence is present to show the foetomaternal placental exchange of insulin<sup>38-40</sup>. In a proposed model of diabetic pregnancy of Pederson it was presumed

that excess maternal glucose can pass through the blood-placental barrier and result in stimulation of endogenous foetal insulin production in the developing foetus rather than direct foetomaternal insulin exchange<sup>41</sup>. There are studies<sup>39,40</sup>, supporting this model where a positive correlation between the cord C peptide along with insulin production and weight of the infant has been shown. This exchange of excess glucose probably can be associated with the insulin resistance of the mother and thereby linking the role of maternal BMI in controlling the level of the excess glucose production and its placental exchange leading to upregulation of foetal insulin secretion and production of bigger babies<sup>42</sup>.

### Conclusion:

It can be concluded that the maternal BMI plays a profound role and can be good indicators of birth weight and development of the fetus<sup>43</sup>. Apart from the risk factors associated with the delivery of a preterm and overweight baby, maternal BMI can be also an important indirect predictor of mothers at risk of delivering abnormal weight babies in developing and poor countries<sup>44,45</sup>. Also, nutrition plays a major role in foetal and maternal well being<sup>42</sup>. Increased risk of negative pregnancy outcomes like pre-term birth, low birth weight, IUGR, small for gestational age (SGA) have shown to be associated with lower maternal pre-BMI<sup>46,47</sup>. However, BMI alone can't be an important anthropometric measurement factor for understanding the growth and development of the foetus. For a more detail understanding of the intra-uterine foetal growth, other important factors like maternal height, weight, metabolic rate, total body water, and foetal body density should be also considered<sup>5,48</sup>.

### References:

1. Gorstein J, Akre J. The use of anthropometry to assess nutritional status. *World Health Stat Q*1988;**41**:48-58.
2. Ay L, Kruithof CJ, Bakker R, Steegers EAP, Witteman JCM, Moll HA, *et al.* Maternal anthropometrics are associated with fetal size in different periods of pregnancy and at birth. The Generation R Study. *Br J Obstet Gynaecol* 2009;**116**:953-63. <https://doi.org/10.1111/j.1471-0528.2009.02143.x>
3. Weiss JL, Malone FD, Emig D, Ball RH, Nyberg DA, Comstock CH, *et al.* Obesity, obstetric complications and cesarean delivery rate-a population-based screening study. *Am J Obstet Gynecol* 2004;**190**:1091-7.
4. Stotland NE, Caughey AB, Breed EM, Escobar GJ. Risk factors and obstetric complications associated with macrosomia. *Int J Gynaecol Obstet* 2004;**87**:220-6.
5. Rasmussen KM, Yaktine AL, editors. Composition and components of gestational weight gain: physiology and metabolism. In: Institute of Medicine (US) and National Research Council (US) Committee to Reexamine IOM Pregnancy Weight Guidelines; Weight Gain during Pregnancy: Reexamining the Guidelines. Washington DC: National Academies Press, 2009.
6. Brown JE, Murtaugh MA, Jacobs DR Jr, Margellos HC. Variation in newborn size according to pregnancy weight change by trimester. *Am J Clin Nutr* 2002;**76**:205-9.
7. Jensen DM, Damm P, Sorensen B, Molsted-Pedersen L, Westergaard JG, Ovesen P, *et al.* Pregnancy outcome and prepregnancy body mass index in 2459 glucose-tolerant Danish women. *Am J Obstet Gynecol* 2003;**189**:239-44.

8. Rode L, Hegaard HK, Kjaergaard H, Moller LF, Tabor A, Ottesen B. Association between maternal weight gain and birth weight. *Obstet Gynecol* 2007;**109**:1309-15.
9. Thorsdottir I, Birgisdottir BE. Different weight gain in women of normal weight before pregnancy: postpartum weight and birth weight. *Obstet Gynecol* 1998;**92**:377-83.
10. Abrams BF, Laros RK Jr. Prepregnancy weight, weight gain, and birth weight. *Am J Obstet Gynecol* 1986;**154**:503-9.
11. Strauss RS, Dietz WH. Low maternal weight gain in the second or third trimester increases the risk for intrauterine growth retardation. *J Nutr* 1999;**129**:988-93.
12. Larciprete G, Valensise H, Vasapollo B, Altomare F, Sorge R, Casalino B, *et al.* Body composition during normal pregnancy: reference ranges. *Acta Diabetol* 2003;**40**:S225-32.
13. Bhattacharya N, Stubblefield PG, editors. Chapter No 6, Anthropometric measurement of the human fetus. Human Fetal Growth and Development, First and Second Trimesters. Switzerland: Springer International Publishing, 2016: 978-3-319-14: 67-83.
14. Chaudhuri A. Embryonic development of human liver and its future implications. In: Bhattacharya N, Stubblefield PG, editors. Human Fetal Growth and Development, First and Second Trimesters. Switzerland: Springer International Publishing, 2016:978-3-319-14:333.
15. Bodey B, Kaiser HE. Development of Hassall's bodies of the thymus in humans and other vertebrates (especially mammals) under physiological and pathological conditions: immunocytochemical, electron microscopic and *in vivo* observations. *In vivo* 1997; **11**:61-85.
16. Dunger DB, Ong KK. Endocrine and metabolic consequences of intrauterine growth retardation. *Endocrinol Metab Clin North Am* 2005;**34**:597-615.
17. Brown JE, Murtaugh MA, Jacobs DR, Margellos HC. Variation in newborn size according to pregnancy weight change by trimester. *Am J Clin Nutr* 2002;**76**:205-9.
18. Abrams B, Selvin S. Maternal weight gain pattern and birth weight. *Obstet Gynecol* 1995;**86**:163-9.
19. Hickey CA, Cliver SP, McNeal SF, Hoffman HJ, Goldenberg RL. Prenatal weight gain patterns and birth weight among nonobese black and white women. *Obstet Gynecol* 1996;**88**:490-6.
20. Li RHJ, Habicht JP. Timing of the influence of maternal nutritional status during pregnancy on fetal growth. *Am J Hum Biol* 1999;**10**:529-39.
21. Scholl TO, Hediger ML, Ances IG, Belsky DH, Salmon RW. Weight gain during pregnancy in adolescence: predictive ability of early weight gain. *Obstet Gynecol* 1990;**75**:948-53.
22. Tan HC, Roberts J, Catov J, Krishnamurthy R, Shypailo R, Bacha F. Mother's pre-pregnancy BMI is an important determinant of adverse cardiometabolic risk in childhood. *Pediatr Diabet* 2015. doi.org/10.1111/pedi.12273 cited by: 22 419-26. PMC. Web 1 May 2018.
23. Wills AK. Maternal and paternal height and BMI and patterns of fetal growth: The Pune Maternal Nutrition Study. *Early Hum Dev* 2010; **86**:9: 535-40. PMC. Web 26 Apr. 2018.
24. Parlee SD, MacDougald OA. Maternal nutrition and risk of obesity in offspring: the Trojan horse of developmental plasticity. *Biochim Biophys Acta* 2014;**1842**:495-506.
25. Goldenberg RL, Davis RO, Cliver SP, Cutter GR, Hoffman HJ, Dubard MB, *et al.* Maternal risk factors and their influence on fetal anthropometric measurements, *Am J Obstet Gynecol* 1993;**168**:1197-205.
26. Zoya T, Farnaz J, Asghar Z, Zohre K, Reza SG, Mitra M. Relationships of the first trimester maternal BMI with new-born anthropometric characteristics and Visfatin Levels throughout pregnancy. *Int J Med Res Health Sci* 2017; **6**: 17-23.
27. Nydjie P, Budd GM, Polansky M. Exploring relationships among maternal BMI, family factors, and concern for child's weight. *J Child Adolesc Psychiatr Nurs* 2010;**23**:4:223-30.

28. Savitz DA, Stein CR, Siega-Riz AM, Herring AH. Gestational weight gain and birth outcome in relation to pre-pregnancy body mass index and ethnicity. *Ann Epidemiol* 2011; **21**:78-85.
29. Dubé MC, Morisset AS, Tchernof A, Weisnagel J. Impact of a mother's weight history on her offspring. *J Obstet Gynaecol Can* 2012;**34**:34-8.
30. Knight B, Shields BM, Turner M, Powell RJ, Yajnik CS, Hattersley AT. Evidence of genetic regulation of fetal longitudinal growth. *Early Hum Dev* 2005;**81**:823-31.
31. Griffiths LJ, Dezateux C, Cole TJ. Differential parental weight and height contributions to offspring birthweight and weight gain in infancy. *Int J Epidemiol* 2007;**36**:104-7.
32. Chen YP, Xiao XM, Li J, Reichetzeder C, Wang ZN, Hocher B. Paternal body mass index (BMI) is associated with offspring intrauterine growth in a gender dependent manner. *PLoS One* 2012;7.5:e36329. PMC.Web.30Apr.2018.
33. Ng SF, Lin RC, Laybutt DR, Barres R, Owens JA, Murriss MJ. Chronic high-fat diet in fathers programs  $\beta$ -cell dysfunction in female rat offspring. *Nature* 2010;**467**:963-6.
34. Luciana Vieira A, Grunebaum A, Alexandra Satty BA, Katherine Sapra J. Fetal ultrasound parameters predict the neonatal body mass index. *Obstet Gynecol* 2014;**123**:101S-2S. doi: 10.1097/01.AOG.0000447447.39668.38.
35. Yalcin BM, Sahin EM, Yalcin E. Which anthropometric measurements is most closely related to elevated blood pressure? *Fam Pract* 2005; **22**:541-7. <https://doi.org/10.1093/fampra/cmi043>
36. Kamath MK; Rao SS, Shenoy RD. Assessment of fetal malnutrition by body mass index and intra uterine growth curves: a comparative study. *Int J Contemporary Pediatr* 2016;**3**:773-7. doi:<http://dx.doi.org/10.18203/2349-3291.ijcp20161415>.
37. Darku ED, Clifford A Jr, Darku MK. Effect of anthropometry and dyslipidaemia on birth weight. *Open Access Library J* 2017; **4**: e3733. <https://doi.org/10.4236/oalib.1103733>.
38. Kalhan SC, Schwartz R, Adam PAJ. Placental barrier to human insulin I125 in Insulin dependent diabetic mothers. *J Clin Endocrinol Metab* 1975;**40**:139.
39. Stanley KP, Fraser RB, Milner M, Bruce C. Cord insulin and C-peptide distribution in an unselected population. *Br J Obstet Gynaecol* 1992;**99**:512-8.
40. Godfrey KM, Hales CN, Osmond C, Barker DJP, Taylor KP. Relation of cord plasma concentrations of proinsulin, 32-33 split proinsulin, insulin and C-peptide to placental weight and the baby's size and proportions at birth. *Early Hum Dev* 1996; **46**:129-40.
41. Retnakaran R1, Ye C, Hanley AJ, Connelly PW, Sermer M, Zinman B, *et al.* Effect of maternal weight, adipokines, glucose intolerance and lipids on infant birth weight among women without gestational diabetes mellitus. *Can Med Assoc J* 2012;**184.12**:1353-60. PMC. Web 1 May 2018.
42. Soltani-KH, Bruce C, Fraser R. Observational study of maternal anthropometry and fetal insulin. *Arch Dis Child Fetal Neonatal Ed* 1999; **81**: F122-4.
43. Li N, Liu E, Guo J, Pan L, Li B, Wang P, *et al.* Maternal prepregnancy body mass index and gestational weight gain on pregnancy outcomes. *PLoS One* 2013; Dec 20, 2013: e82310. web 1 May 2018.
44. Golditch IM, Kirkman K. The large fetus, management and outcome. *Obstet Gynecol* 1978; **52**:26-30.
45. Langer O, Berkus MD, Huff RW, Samueloof A. Shoulder dystocia: should the fetus weighing greater than or equal to 4000 grams be delivered by cesarean section? *Am J Obstet Gynecol* 1991;**165**:831-7.
46. Liu Y, Dai W, Dai X, Li Z. Pre-pregnancy body mass index and gestational weight gain with the outcome of pregnancy: a 13-year study of 292,568 cases in China. *Arch Gynecol Obstet* 2012;**286**: 905-11.
47. Li N, Liu E, Guo J, Pan L, Li B, Wang P, *et al.* Maternal pre-pregnancy body mass index and gestational weight gain on offspring overweight in early infancy. *PLoS One* 2013;**8**: e77809.

48. Parsons TJ, Power C, Manor O. Foetal and early life growth and body mass index from birth to early adulthood in 1958 British cohort: longitudinal study. *BMJ* 2001; **323**:1331-5.